

INDIVIDUAL ERGODIC THEOREMS IN NONCOMMUTATIVE SYMMETRIC SPACES

VLADIMIR CHILIN AND SEMYON LITVINOV

ABSTRACT. It is known that, for a positive Dunford-Schwartz operator in a noncommutative L^p -space, $1 \leq p < \infty$ or, more generally, in a noncommutative Orlicz space with order continuous norm, the corresponding ergodic averages converge bilaterally almost uniformly. We show that these averages converge bilaterally almost uniformly in each noncommutative symmetric space E such that $\mu_t(x) \rightarrow 0$ as $t \rightarrow 0$ for every $x \in E$, where $\mu_t(x)$ is a non-increasing rearrangement of x . In particular, these averages converge bilaterally almost uniformly in all noncommutative symmetric spaces with order continuous norm.

1. PRELIMINARIES

Let $\mathcal{M}$ be a semifinite von Neumann algebra equipped with a faithful normal semifinite trace τ . Let $\mathcal{P}(\mathcal{M})$ stand for the set of projections in $\mathcal{M}$. If $\mathbf{1}$ is the identity of $\mathcal{M}$ and $e \in \mathcal{P}(\mathcal{M})$, we write $e^\perp = \mathbf{1} - e$. Denote by $L^0 = L^0(\mathcal{M}) = L^0(\mathcal{M}, \tau)$ the $*$ -algebra of τ -measurable operators affiliated with $\mathcal{M}$. Let $\|\cdot\|_\infty$ be the uniform norm in $\mathcal{M}$. Equipped with the *measure topology* given by the system

$$V(\epsilon, \delta) = \{x \in L^0 : \|xe\|_\infty \leq \delta \text{ for some } e \in \mathcal{P}(\mathcal{M}) \text{ with } \tau(e^\perp) \leq \epsilon\},$$

$\epsilon > 0, \delta > 0$, L^0 is a complete metrizable topological $*$ -algebra [14].

Let $x \in L^0$, and let $\{e_\lambda\}_{\lambda \geq 0}$ be the spectral family of projections for the absolute value $|x|$ of x . If $t > 0$, then the t -th *generalized singular number of x* (a *non-increasing rearrangement of x*) (see [9]) is defined as

$$\mu_t(x) = \inf\{\lambda > 0 : \tau(e_\lambda^\perp) \leq t\}.$$

A Banach space $(E, \|\cdot\|_E) \subset L^0$ is called *fully symmetric* if the conditions

$$x \in E, y \in L^0, \int_0^s \mu_t(y) dt \leq \int_0^s \mu_t(x) dt \text{ for all } s > 0 \text{ (writing } y \prec\prec x)$$

imply that $y \in E$ and $\|y\|_E \leq \|x\|_E$. It is known [6] that if $(E, \|\cdot\|_E)$ is a fully symmetric space, $x_n, x \in E$, and $\|x - x_n\|_E \rightarrow 0$, then $x_n \rightarrow x$ in measure. A fully symmetric space $(E, \|\cdot\|_E)$ is said to possess *Fatou property* if conditions

$$x_\alpha \in E^+, x_\alpha \leq x_\beta \text{ for } \alpha \leq \beta, \text{ and } \sup_\alpha \|x_\alpha\|_E < \infty$$

Date: April 1, 2016.

2010 Mathematics Subject Classification. 47A35(primary), 46L52(secondary).

Key words and phrases. Semifinite von Neumann algebra, Dunford-Schwartz operator, non-commutative symmetric space, individual ergodic theorem.

imply that there exists $x = \sup_{\alpha} x_{\alpha} \in E$ and $\|x\|_E = \sup_{\alpha} \|x_{\alpha}\|_E$. A space $(E, \|\cdot\|_E)$ is said to have *order continuous norm* if $\|x_{\alpha}\|_E \downarrow 0$ whenever $x_{\alpha} \in E$ and $x_{\alpha} \downarrow 0$.

Let $L^0(0, \infty)$ be the linear space of all (equivalence classes of) almost everywhere finite complex-valued Lebesgue measurable functions on the interval $(0, \infty)$. We identify $L^{\infty}(0, \infty)$ with the commutative von Neumann algebra acting on the Hilbert space $L^2(0, \infty)$ via multiplication by the elements from $L^{\infty}(0, \infty)$ with the trace given by the integration with respect to Lebesgue measure. A Banach space $E \subset L^0(0, \infty)$ is called *fully symmetric function space* on $(0, \infty)$ if the condition above holds with respect to the von Neumann algebra $L^{\infty}(0, \infty)$.

Let $E = (E(0, \infty), \|\cdot\|_E)$ be a fully symmetric function space. For each $s > 0$ let $D_s : E \rightarrow E$ be the bounded linear operator given by $D_s(f)(t) = f(t/s)$, $t > 0$. The *Boyd indices* p_E and q_E are defined as

$$p_E = \lim_{s \rightarrow \infty} \frac{\log s}{\log \|D_s\|_E}, \quad q_E = \lim_{s \rightarrow +0} \frac{\log s}{\log \|D_s\|_E}.$$

It is known that $1 \leq p_E \leq q_E \leq \infty$ [13, II, Ch.2, Proposition 2.b.2]. A fully symmetric function space is said to have *non-trivial Boyd indices* if $1 < p_E$ and $q_E < \infty$. For example, the space $L^p(0, \infty)$, $1 < p < \infty$, have non-trivial Boyd indices:

$$p_{L^p(0, \infty)} = q_{L^p(0, \infty)} = p$$

[1, Ch.4, §4, Theorem 4.3].

A Banach lattice $(E, \|\cdot\|_E)$ is called *q-concave*, $1 \leq q < \infty$, if there exists a constant $M > 0$ such that

$$\left(\sum_{i=1}^n \|x_i\|^q \right)^{\frac{1}{q}} \leq M \left\| \left(\sum_{i=1}^n |x_i|^q \right)^{\frac{1}{q}} \right\|_E$$

for every finite set $\{x_i\}_{i=1}^n \subset E$. Let $E = (E(0, \infty), \|\cdot\|_E)$ be a fully symmetric function space, and let E be *q-concave* for same $1 \leq q < \infty$. Then there does not exist a sublattice of E isomorphic to l^{∞} , hence the norm $\|\cdot\|_E$ is order continuous [16, Corollary 2.4.3].

If $E(0, \infty)$ is a fully symmetric function space, define

$$E(\mathcal{M}) = E(\mathcal{M}, \tau) = \{x \in L^0 : \mu_t(x) \in E\}$$

and set

$$\|x\|_{E(\mathcal{M})} = \|\mu_t(x)\|_E, \quad x \in E(\mathcal{M}).$$

It is shown in [12] that $(E(\mathcal{M}), \|\cdot\|_{E(\mathcal{M})})$ is a fully symmetric space. If $1 \leq p < \infty$ and $E = L_p(0, \infty)$, the space $(E(\mathcal{M}), \|\cdot\|_{E(\mathcal{M})})$ coincides with the noncommutative L^p -space $L^p = L^p(\mathcal{M}) = (L^p(\mathcal{M}, \tau), \|\cdot\|_p)$ because

$$\|x\|_p = \left(\int_0^{\infty} \mu_t^p(x) dt \right)^{1/p} = \|x\|_{L^p(\mathcal{M})}$$

(see [17, Proposition 2.4]). In addition, $L^{\infty}(\mathcal{M}) = \mathcal{M}$ and

$$(L^1 \cap L^{\infty})(\mathcal{M}) = L^1(\mathcal{M}) \cap \mathcal{M}, \quad \|x\|_{L^1 \cap L^{\infty}} = \max\{\|x\|_1, \|x\|_{\mathcal{M}}\},$$

$$(L^1 + L^{\infty})(\mathcal{M}) = L^1(\mathcal{M}) + \mathcal{M},$$

$$\|x\|_{L^1+L^\infty} = \inf\{\|x\|_1 + \|y\|_{\mathcal{M}} : x \in L^1(\mathcal{M}), y \in \mathcal{M}\} = \int_0^1 \mu_t(x) dt$$

(see [6, Proposition 2.5]). (For a comprehensive review of noncommutative L^p -spaces, see [17, 15].)

Since for a fully symmetric function space $E(0, \infty)$,

$$L^1(0, \infty) \cap L^\infty(0, \infty) \subset E \subset L^1(0, \infty) + L^\infty(0, \infty)$$

with continuous embeddings [11, Ch.II, §4, Theorem 4.1], we also have

$$L^1(\mathcal{M}) \cap \mathcal{M} \subset E(\mathcal{M}, \tau) \subset L^1(\mathcal{M}) + \mathcal{M},$$

with continuous embeddings.

Denote

$$L_\tau^0 = \{x \in L^0(\mathcal{M}) : \mu_t(x) \rightarrow 0 \text{ as } t \rightarrow 0\}.$$

Observe that L_τ^0 is a linear subspace of L^0 which is solid in the sense that if $x \in L_\tau^0$ and if $y \in L^0$ and $\mu_t(y) \leq \mu_t(x)$, then also $y \in L_\tau^0$. It is clear that $x \in L_\tau^0$ if and only if $\tau(\chi_{(\lambda, \infty)}(|x|)) < \infty$ for all $\lambda > 0$; moreover, $L^1 \cap \mathcal{M} \subset L_\tau^0$.

Define

$$\mathcal{R}_\tau = \{x \in L^1 + \mathcal{M} : \mu_t(x) \rightarrow 0 \text{ as } t \rightarrow 0\}.$$

It follows from the next proposition that $(\mathcal{R}_\tau, \|\cdot\|_{L^1+\mathcal{M}})$ is a Banach space.

Proposition 1.1. [6, Proposition 2.7] $\mathcal{R}_\tau$ is the closure of $L^1 \cap \mathcal{M}$ in $L^1 + \mathcal{M}$.

Note that definitions of L_τ^0 and $\|\cdot\|_{L^1+L^\infty}$ yield that if

$$x \in \mathcal{R}_\tau, y \in L^1 + L^\infty, \text{ and } \mu_t(y) \leq \mu_t(x),$$

then $y \in \mathcal{R}_\tau$ and $\|y\|_{L^1+\mathcal{M}} \leq \|x\|_{L^1+\mathcal{M}}$. Therefore $(\mathcal{R}_\tau, \|\cdot\|_{L^1+\mathcal{M}})$ is a noncommutative symmetric space [6, §2].

2. INDIVIDUAL ERGODIC THEOREMS IN $\mathcal{R}_\tau$

A linear operator $T : L^1 + \mathcal{M} \rightarrow L^1 + \mathcal{M}$ is called a *Dunford-Schwartz operator* (writing $T \in DS$) if

$$\|T(x)\|_1 \leq \|x\|_1 \quad \forall x \in L^1 \quad \text{and} \quad \|T(x)\|_\infty \leq \|x\|_\infty \quad \forall x \in \mathcal{M}.$$

If a Dunford-Schwartz operator T is positive, that is, $T(x) \geq 0$ whenever $x \geq 0$, we shall write $T \in DS^+$.

If $T \in DS$, then

$$(1) \quad \|T\|_{L^1+\mathcal{M} \rightarrow L^1+\mathcal{M}} \leq 1$$

(see, for example, [6, §4]) and, in addition, $Tx \prec\prec x$ for all $x \in L^1 + \mathcal{M}$ [6, Theorem 4.7]. Hence

$$(2) \quad T(\mathcal{R}_\tau) \subset \mathcal{R}_\tau.$$

Given $T \in DS$ and $x \in L^1 + L^\infty$, denote

$$(3) \quad A_n(x) = A_n(T, x) = \frac{1}{n} \sum_{k=0}^{n-1} T^k(x), \quad n = 1, 2, \dots,$$

the corresponding ergodic averages of the operator x .

Note that, by (1) and (2), for any $T \in DS$ its restriction on $\mathcal{R}_\tau$ is a linear contraction (also denoted by T).

Definition 2.1. A sequence $\{x_n\} \subset L^0$ is said to converge to $x \in L^0$ *bilaterally almost uniformly (b.a.u.)* if for every $\epsilon > 0$ there exists such a projection $e \in \mathcal{P}(\mathcal{M})$ that $\tau(e^\perp) \leq \epsilon$ and $\|e(x - x_n)e\|_\infty \rightarrow 0$.

The following groundbreaking result was established in [18] as a corollary of a noncommutative maximal ergodic inequality [18, Theorem 1] (for the assumption $T \in DS^+$, see [4, Remark 1.2] and [10, Lemma 1.1]).

Theorem 2.1. *Let $T \in DS^+$ and $x \in L^1$. Then the averages (3) converge b.a.u. to some $\hat{x} \in L^1$.*

The next result is an extension of Theorem 2.1 to $\mathcal{R}_\tau$.

Theorem 2.2. *Let $T \in DS^+$ and $x \in \mathcal{R}_\tau$. Then the averages (3) converge b.a.u. to some $\hat{x} \in L^1 + \mathcal{M}$.*

Proof. Without loss of generality assume that $x \geq 0$. Let $\{e_\lambda\}_{\lambda \geq 0}$ be the spectral family of x . Given $k \in \mathbb{N}$, denote $x_k = \int_{1/k}^\infty \lambda de_\lambda$ and $y_k = \int_0^{1/k} \lambda de_\lambda$. Then $0 \leq y_k \leq \frac{1}{k} \cdot \mathbf{1}$, $x_k \in L^1$, and $x = x_k + y_k$ for all k .

Fix $\epsilon > 0$. By Theorem 2.1, $A_n(x_k) \rightarrow \hat{x}_k \in L^1$ b.a.u. for each k . Therefore there exists $e_k \in \mathcal{P}(\mathcal{M})$ such that $\tau(e_k^\perp) \leq \frac{\epsilon}{2^k}$ and $\|e_k(A_n(x_k) - \hat{x}_k)e_k\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.

Then it follows that

$$\|e_k(A_n(x_k) - A_m(x_k))e_k\|_\infty < \frac{1}{k} \quad \text{for all } m, n \geq N(k).$$

Since $\|y_k\|_\infty \leq \frac{1}{k}$, we have

$$\begin{aligned} & \|e_k(A_n(x) - A_m(x))e_k\|_\infty \leq \\ & \leq \|e_k(A_n(x_k) - A_m(x_k))e_k\|_\infty + \|e_k(A_n(y_k) - A_m(y_k))e_k\|_\infty < \\ & < \frac{1}{k} + \|e_k A_n(y_k)e_k\|_\infty + \|e_k A_m(y_k)e_k\|_\infty \leq \frac{3}{k}. \end{aligned}$$

for each k and all $m, n \geq N(k)$.

Let $e = \bigwedge_{k \geq 1} e_k$. Then $\tau(e^\perp) \leq \sum_{k=1}^\infty \tau(e_k^\perp) \leq \epsilon$ and

$$\|e(A_n(x) - A_m(x))e\|_\infty < \frac{3}{k}$$

for all $m, n \geq N(k)$. This means that $\{A_n(x)\}$ is Cauchy with respect to b.a.u. convergence. Then, by [3, Theorem 2.3], we conclude that the sequence $\{A_n(x)\}$ converges b.a.u. to some $\hat{x} \in L^0$.

Since $L^1 + \mathcal{M}$ satisfies Fatou property [8, §4], its unit ball is closed in the measure topology [7, Theorem 4.1], and (1) implies that $\hat{x} \in L^1 + \mathcal{M}$. $\square$

Let $\mathbb{C}_1 = \{z \in \mathbb{C} : |z| = 1\}$ be the unit circle in $\mathbb{C}$. A function $P : \mathbb{Z} \rightarrow \mathbb{C}$ is said to be a *trigonometric polynomial* if $P(k) = \sum_{j=1}^s z_j \lambda_j^k$, $k \in \mathbb{Z}$, for some $s \in \mathbb{N}$, $\{z_j\}_1^s \subset \mathbb{C}$, and $\{\lambda_j\}_1^s \subset \mathbb{C}_1$. A sequence $\{\beta_k\}_{k=0}^\infty \subset \mathbb{C}$ is called a *bounded Besicovitch sequence* if

- (i) $|\beta_k| \leq C < \infty$ for all k ;
- (ii) for every $\epsilon > 0$ there exists a trigonometric polynomial P such that

$$\limsup_n \frac{1}{n+1} \sum_{k=0}^n |\beta_k - P(k)| < \epsilon.$$

The following theorem was established in [3].

Theorem 2.3. *Assume that $\mathcal{M}$ has a separable predual. Let $T \in DS^+$ and let $\{\beta_k\}$ be a Besicovitch sequence. Then for every $x \in L^1(\mathcal{M})$ the averages*

$$(4) \quad \frac{1}{n} \sum_{k=0}^{n-1} \beta_k T^k(x)$$

converge b.a.u. to some $\hat{x} \in L^1$.

In view of Theorem 2.3 and since a sequence $\{\beta_k\}$ in question is bounded, the proof of Theorem 2.2 can be carried out for the averages (4), and we obtain the following.

Theorem 2.4. *Let $\mathcal{M}$, T , $\{\beta_k\}$ be as in Theorem 2.3. Then for every $x \in \mathcal{R}_\tau$ the averages (4) converge to some $\hat{x} \in L^1 + \mathcal{M}$.*

3. APPLICATIONS OF THEOREM 2.2

Let us present some examples of noncommutative symmetric spaces for which Dunford-Schwartz individual ergodic theorem is valid.

1. Let Φ be an Orlicz function, $L^\Phi = L^\Phi(\mathcal{M}, \tau)$ the corresponding noncommutative Orlicz space, $\|\cdot\|_\Phi$ the Luxemburg norm in L^Φ (see [5]). Since $(L^\Phi(0, \infty), \|\cdot\|_\Phi)$ has Fatou property, the space $(L^\Phi(\mathcal{M}, \tau), \|\cdot\|_\Phi)$ has it as well [6, §5]. As shown in the proof of [5, Proposition 2.1], $L^\Phi \subset \mathcal{R}_\tau$. It follows then from Theorem 2.2 that we have

Theorem 3.1. *Let $T \in DS^+$ and $x \in L^\Phi$. Then the averages (3) converge b.a.u. to some $\hat{x} \in L^\Phi$.*

2. Let $(E(\mathcal{M}), \|\cdot\|_{E(\mathcal{M})})$ be a noncommutative fully symmetric space with order continuous norm. Then $\tau(\{|x| > \lambda\}) < \infty$ for all $x \in E(\mathcal{M})$ and $\lambda > 0$, so $E(\mathcal{M}) \subset \mathcal{R}_\tau$. Thus we have

Theorem 3.2. *Let $(E(\mathcal{M}), \|\cdot\|_{E(\mathcal{M})})$ be noncommutative fully symmetric space with order continuous norm. Let $T \in DS^+$ and $x \in E(\mathcal{M})$. Then the averages (3) converge b.a.u. to some $\hat{x} \in E(\mathcal{M})$.*

3. Let φ be an increasing continuous concave function on $[0, \infty)$ with $\varphi(0) = 0$ and $\varphi(\infty) = \infty$, and let $\Lambda_\varphi(\mathcal{M})$ be the corresponding noncommutative Lorentz space (see, for example, [2]). Since $\Lambda_\varphi(0, \infty)$ is a fully symmetric space with order continuous norm and Fatou property [11, Ch. II, §5], $\Lambda_\varphi(\mathcal{M})$ also has order continuous norm and Fatou property [6, Proposition 3.6 and §5]. Then we have

Theorem 3.3. *Let $T \in DS^+$ and $x \in \Lambda_\varphi(\mathcal{M})$. Then the averages (3) converge b.a.u. to some $\hat{x} \in \Lambda_\varphi(\mathcal{M})$.*

4. Let $E = (E(0, \infty), \|\cdot\|_{E(0, \infty)})$ be a fully symmetric function space, and let E be q -concave for some $1 \leq q < \infty$. Then E , hence $(E(\mathcal{M}), \|\cdot\|_{E(\mathcal{M})})$, has order continuous norm. Therefore we have the following.

Theorem 3.4. *Let $E = (E(0, \infty), \|\cdot\|_{E(0, \infty)})$ be a fully symmetric function space. Assume that E is q -concave for some $1 \leq q < \infty$. Let $T \in DS^+$ and $x \in E(\mathcal{M})$. Then the averages (3) converge b.a.u. to some $\hat{x} \in L^1 + \mathcal{M}$. In addition, if E has Fatou property, then $\hat{x} \in E(\mathcal{M})$.*

5. Let $E = (E(0, \infty), \|\cdot\|_{E(0, \infty)})$ be a fully symmetric function space. Assume that E has non-trivial Boyd indices. According to [13, II, Ch.2, Proposition 2.b.3], there exist such $1 < p, q < \infty$ that the space E is intermediate for the Banach couple $(L^p(0, \infty), L^q(0, \infty))$. Since

$$(L^p + L^q)(\mathcal{M}) = L^p(\mathcal{M}) + L^q(\mathcal{M})$$

(see [6, Proposition 3.1]), we have

$$E(\mathcal{M}) \subset L^p(\mathcal{M}) + L^q(\mathcal{M}) \subset \mathcal{R}_\tau.$$

Then we also have (cf. [4, Theorem 4.2])

Theorem 3.5. *Let $E = (E(0, \infty), \|\cdot\|_{E(0, \infty)})$ be a fully symmetric function space, and let E have non-trivial Boyd indices. Given $T \in DS^+$ and $x \in E(\mathcal{M})$, the averages (3) converge b.a.u. to some $\hat{x} \in L^1 + \mathcal{M}$. In addition, if E has Fatou property, then $\hat{x} \in E(\mathcal{M})$.*

Remark 3.1. It is clear that if one assumes that $\mathcal{M}$ has a separable predual, Theorems 3.1 - 3.5 remain valid for the Besicovitch weighted averages (4).

REFERENCES

- [1] C. Bennett, R. Sharpley, *Interpolation of Operators*, Academic Press Inc. (London) LTD, 1988.
- [2] V. Chilin, F. Sukochev, Weak convergence in non-commutative symmetric spaces, *J. Operator Theory*, **31** (1994), 35-55.
- [3] V. Chilin, S. Litvinov, and A. Skalski, A few remarks in non-commutative ergodic theory, *J. Operator Theory*, **53**(2) (2005), 331-350.
- [4] V. Chilin, S. Litvinov, Ergodic theorems in fully symmetric spaces of τ -measurable operators, *Studia Math.*, **288**(2) (2015), 177-195.
- [5] V. Chilin, S. Litvinov, Individual ergodic theorems in noncommutative Orlicz spaces, *Positivity*, DOI 10.1007/s11117-016-0402-8.
- [6] P. G. Dodds, T. K. Dodds, and B. Pagter, Noncommutative Köthe duality, *Trans. Amer. Math. Soc.*, **339**(2) (1993), 717-750.
- [7] P. G. Dodds, T. K. Dodds, F. A. Sukochev, and O. Ye. Tikhonov, A Non-commutative Yosida-Hewitt theorem and convex sets of measurable operators closed locally in measure, *Positivity*, **9** (2005), 457-484.
- [8] P. G. Dodds, B. Pagter, The non-commutative Yosida-Hewitt decomposition revisited, *Trans. Amer. Math. Soc.*, **364**(12) (2012), 6425-6457.
- [9] T. Fack, H. Kosaki, Generalized s -numbers of τ -measurable operators, *Pacific. J. Math.*, **123** (1986), 269-300.
- [10] M. Junge, Q. Xu, Noncommutative maximal ergodic theorems, *J. Amer. Math. Soc.*, **20**(2) (2007), 385-439.
- [11] S. G. Krein, Ju. I. Petunin, and E. M. Semenov, *Interpolation of Linear Operators*, Translations of Mathematical Monographs, Amer. Math. Soc., **54**, 1982.
- [12] N. J. Kalton, F. A. Sukochev, Symmetric norms and spaces of operators, *J. Reine Angew. Math.*, **621** (2008), 81-121.
- [13] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces I-II*, Springer-Verlag, Berlin Heidelberg New York, 1977.
- [14] E. Nelson, Notes on non-commutative integration, *J. Funct. Anal.*, **15** (1974), 103-116.
- [15] G. Pisier, Q. Xu, Noncommutative L^p -spaces, in *Handbook of the geometry of Banach spaces*, **2** (2003), 1459-1517.
- [16] Peter Meyer-Niebel, *Banach Lattices*, Springer-Verlag, New York Berlin Heidelberg, 1991.
- [17] F. J. Yeadon, Non-commutative L^p -spaces, *Math. Proc. Camb. Phil. Soc.*, **77** (1975), 91-102.
- [18] F. J. Yeadon, Ergodic theorems for semifinite von Neumann algebras. I, *J. London Math. Soc.*, **16**(2) (1977), 326-332.

THE NATIONAL UNIVERSITY OF UZBEKISTAN
E-mail address: `vladimirchil@gmail.com`

PENNSYLVANIA STATE UNIVERSITY HAZLETON
E-mail address: `snl2@psu.edu`